





ALLTogether

		IR-High Maintenance with VCR-Dexa pulses																											
Week	37							38							39							40							
Day	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	etc
6-MP 75 mg/m <sup>2</sup>	[Blue bar]																												
MTX 20 mg/m <sup>2</sup>	[Yellow bar]							[Yellow bar]							[Yellow bar]							[Yellow bar]							
IT according to CNS/age	*																												
VCR 1.5 mg/m <sup>2</sup>	[Red bar]																												
Dexa 6 mg/m <sup>2</sup>	[Light blue bar]																												
*Every 3 <sup>rd</sup> cycle (every 12 weeks) throughout Maintenance																													

ELLER hvis randomiseret til InO i R3

		IR-High Inotuzumab experimental arm																																								
Week	37							38							39							40							41							42						
Day	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294
InO 0.5 mg/m <sup>2</sup>	[Purple bar]							[Purple bar]							[Purple bar]							[Purple bar]							[Purple bar]													
Cycle 1														Cycle 2																												

		IR-High Maintenance with VCR-Dexa pulses post InO																											
Week	43							44							45							46							
Day	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	etc
6-MP 75 mg/m <sup>2</sup>	[Blue bar]																												
MTX 20 mg/m <sup>2</sup>	[Yellow bar]							[Yellow bar]							[Yellow bar]							[Yellow bar]							
IT according to CNS/age	*																												
VCR 1.5 mg/m <sup>2</sup>	[Red bar]																												
Dexa 6 mg/m <sup>2</sup>	[Light blue bar]																												
*Every 3 <sup>rd</sup> cycle (every 12 weeks), throughout Maintenance																													

Eligible patients:

**This part of the protocol applies to**

- Patients who have received the experimental Inotuzumab block and should continue Standard Maintenance.

### Standard Risk

	Induction (NCI High Risk/T-cell)																												
Week	1							2							3							4							
Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
Dexa 6 mg/m <sup>2</sup>																													
VCR 1.5 mg/m <sup>2</sup>																													
DNR 25 mg/m <sup>2</sup>																													
PEG-Asp 1500/1000 IU/m <sup>2</sup>			1															2											
ITT, according to age								*																					
	*CNS2/3 and TLP+																												
BM	x														x														

TREATMENT PHASE	DURATION	DETAILS OF TREATMENTS/RANDOMISATIONS
Consolidation 1	6 weeks (weeks 5-10)	
Consolidation 2	7 weeks (weeks 11-17)	
Delayed Intensification	7 weeks (weeks 18-24)	Randomisation 1. With and without Anthracycline (Doxorubicin)
Maintenance	Continues for exactly 2 years from the start of consolidation 1	

